

J. Theodore Schwartz, Jr M.D.  
Webster Orthopedic Medical Group  
5801 Norris Canyon Rd, Suite 210  
San Ramon, CA 94583  
(925) 355-7350

## **After Surgery**

### **Instructions after Trigger Finger Release**

**General Information:** During surgery a local anesthetic was injected into your finger, so after surgery your finger may be numb from 4 to 48 hours.

- Pain Management** Take the prescribed pain medication for pain. Follow the direction on the package. If there are any question regarding the pain medication, please call our office. If you wish, you may take over-the-counter pain medication such as acetaminophen (Tylenol), ibuprofen (Advil, Nuprin, or Motrin), naproxen (Aleve) instead. It is recommended to take pain medication with food.
- Icing and Elevation** To decrease swelling, it is helpful to elevate your hand above your heart and apply an ice pack to your palm for the first 24 hours. You may use a sling if you wish for the first 24 hours after surgery, but remember a sling will typically keep your hand below your heart.
- Dressings** Keep the dressing clean and dry. If the Ace bandage feels tight, unwrap the Ace bandage and rewrap it so it feels comfortable. You can remove the dressing three days after surgery and place a Band-aid over the incision. Do not apply bacitracin to the wound unless specifically instructed.
- Activities** The day following surgery, you may perform light activities with the finger. You can perform anything that does not cause pain at the incision or in the finger. You are encouraged to move your fingers as soon as is comfortable so that they do not become stiff.
- Follow-Up** You should have a follow-up appointment for approximately 10 days after surgery. If you do not have a follow-up appointment, please call the office to arrange one.

If something seems wrong or out of the ordinary please contact the office at 925-355-7350